



Across

- Degenerative joint disease
- Carpal tunnel symptoms are usually reversible after the _____ is incised.
- _____ is an abnormal thickening of the tissue beneath the skin. This thickening occurs in the palm and can extend into the fingers.
- Carpal tunnel release may be completed _____ or by open incision.
- The four basic instruments used in hand surgery are a knife, small forceps, dissecting scissors, and _____.

Down

- The medical term for "trigger finger"
- Number of muscles that move the thumb
- Carpal tunnel syndrome is caused by thickened _____, trauma, or aberrant muscles.
- The reconstructive procedure of toe-to-hand transfer involves surgical removal of a single toe or multiple toes and _____ of the vessels of the toes to those on the hand to restore finger and thumb functions.
- A cystic lesion arising from a joint capsule or tendon sheath containing clear fluid
- Arteries and veins should be reconnected without tension. If this cannot be done the hand surgeon must take out a piece of vein from another place in the patient's body and use it to reconstruct the vein in the hand. This process is called a _____.
- Ganglions are most common on the _____ of the wrist or palm of the hand.



Clues taken from:
 Surgery Encyclopedia
<http://www.surgeryencyclopedia.com/FiLa/HandSurgery.html#ixzz3ZxKjyNcr>
 American Society for Hand Surgery
<http://www.assh.org/handcare/>