

Across


- _____ hernias occur due to a weakness, gap, or opening in the muscles or tendons of the upper abdominal wall, on a line between the breast bone and the navel or umbilicus.
- A hernia that appears in the abdomen at the site of a previous surgery is known as a ventral or _____ hernia.
- These hernias represent approximately two thirds of adult hernias and are much more common in men than in women.
- _____ repair techniques use a piece of mesh to bridge the hernia defect.
- Femoral hernias are prone to develop incarceration (tissues of the hernia are trapped) or _____ (the tissue is trapped and blood supply cut off).
- These hernias may occur in infants at or just after birth and may resolve by three or four years of age.
- Children's hernias are almost always _____, opposed to acquired.

Down

- Laparoscopic Tension Free Repair surgery is called _____ hernia repair because the hernia is being repaired from behind the abdominal wall.
- When inguinal hernias are repaired with a technique called _____, recurrence rates may be higher than 15%, which means that the hernia may reappear in time.
- _____ hernias are slightly different, they are a weakness or opening in the diaphragm which can lead to heartburn, pain, and erosion of the esophagus.
- An indirect inguinal hernia occurs through the natural weakness in the internal _____.
- Sterile, woven material made from a synthetic plastic-like material.
- A direct inguinal hernia is a result of weakness in the floor of the _____ and is more likely to develop in men over the age of 40.
- These hernias appear just below the groin crease and are usually the result of pregnancy and childbirth.

Reference:

www.herniasolutions.com



www.herniaguide.com