



Across

- PERITONEALLAVAGE**—The trauma patient who is hypotensive and has left upper quadrant pain can require _____ to determine abdominal bleeding
- HEMOTHORAX**—A collection of blood in the pleural space can be caused by either blunt or penetrating trauma. This collection of blood is known as _____.
- COMPOUND**—This type of fracture is considered a surgical emergency
- BURRHOLES**—A patient who presents status post fall that resulted in subdural hematoma. The procedure used in treatment is :
- LUMBAR**—Cauda Equina is a spinal injury that occurs in what region of the spine?
- III**—The LeFort _____ involves the zygomatic arch
- HEARINGLOSS**—Close proximity to an explosion can result in _____
- PENETRATING**—This type of chest wound is the result of a projectile or knife.
- RAINEY**—Hemostasis of scalp edges during craniotomy requires a _____ clip(s).
- CRANIOPLASTY**—The procedure to correct skull defects is known as _____

Down

- LOGROLLED**—Patients with suspected spinal injuries should be _____ when positioning
- GENITOURINARY**—As a result of _____ war injuries, the military has started providing Kevlar underpants as added protection for the troops
- HEMMORHAGE**—An intraoperative complication with splenectomy procedures is _____
- QUADRIPLEGIA**—Injuries that occur in the cervical region result in _____
- AMPUTATION**—Traumatic limb injury can result in: _____
- ENUCLEATION**—The correct procedure name for removing a damaged eye is _____

