



Across

4. Gas filling the abdomen during laparoscopy results in _____ being created
6. This syndrome is one cause of intestinal failure. (2 words)
8. Chronic exposure to gastric acid in the esophagus can result in _____ esophagus.
9. _____ are used to aid in inserting instruments into the abdomen during laparoscopic cases.
12. _____ is the protrusion of bowel from an open wound
13. The GI tract is also known as the _____ canal.
15. Separation of wound edges is known as:

Down

1. Cell salvage is also referred to as:
2. In determining the surgical approach for esophagectomy, a surgeon must consider the _____ and location of the tumor.
3. Cross contamination of malignant cells is also referred to as:
5. If an appendectomy is performed as an open technique, the surgeon will commonly use the _____ incision
7. This instrument is utilized to visualize the abdominal cavity
10. _____ is the double layer of fatty peritoneum that is attached to the greater curve of the stomach.
11. Weight reduction surgery is also considered _____ surgery.
14. To help decrease the risk of _____ in GI Surgery patients, intermittent pneumatic compression devices are used

To complete this activity, the participants may need to access and read the "Gastrointestinal Surgery" chapter within *Alexander's Care of the Surgical Patient*. Rothrock, J.C. (Ed.). (2015). St. Louis, MO: Elsevier., p. 295-349.