



To complete this activity, the participant will need to access and read McEwen, D.R. (2019). Pediatric Surgery. In J.C. Rothrock (Ed.). *Alexander's Care of the Patient in Surgery* (pp. 999 - 1066). St. Louis, MO: Elsevier.

**Across**

- 2. Most common cause of small bowel obstruction in children.
- 3. Most common cause of gastric outlet obstruction in children. (2 words)
- 5. Excision of foreskin of the glans penis.
- 7. \_\_\_\_\_ is the most serious risk of foreign body ingestion.
- 12. Surgical placement of the testicle in the scrotal sac.
- 14. Condition in which the ductus arteriosus fails to close within 24 - 48 hours after birth.
- 15. Third most common childhood cancer.
- 16. \_\_\_\_\_ tumor is also known as nephroblastoma.
- 17. Infants are more prone to \_\_\_\_\_ as a result of immature renal function and limited kidney function.

**Down**

- 1. The predominate determinant of cardiac output in infants and children. (2 words)
- 3. An obvious defect of the sternum giving the appearance of a funnel chest. (2 words)
- 4. Fetal surgery involves at a minimum \_\_\_\_\_ (number) patients.
- 6. A temporary or permanent channel from the gastric lumen to the skin.
- 8. Surgical procedure for children and infants that experience severe GERD.
- 9. Urethral opening on the dorsum of the penis.
- 10. Core body temperature lower than 37 degrees Celsius.
- 11. Preoperative sedation is generally accomplished using \_\_\_\_\_ which can be administered orally.
- 13. Craniosynostosis that involves only one suture line is classified as \_\_\_\_\_.

\* Content reviewed by the CCI Nursing Education Department for alignment with clinical practice standards. CCI does not require, recommend, or endorse specific training programs in specialized practice areas for any of its exams. This is an example of a future points activity for recertification in collaboration with CCI.