



Visit www.mhaus.org for reference.

Across

- 2. Drug used to treat hyperkalemia. (2 words)
- 5. Temperature above 38 degrees Celsius.
- 7. If the patient has had an MH episode, they should be monitored in an ICU setting for _____ hours. (2 words)
- 9. The only diluent that should be used for Dantrolene. (2 words)
- 12. In patients with this form of muscular dystrophy, succinylcholine should be avoided.
- 13. Muscles may become _____ during an MH crisis.
- 15. The commonly known anesthetic trigger for Malignant Hyperthermia.

Down

- 1. The standard for MH diagnostic testing. (acronym)
- 3. A test used to determine the degree of metabolic acidosis in a patient during MH crisis. (3 words)
- 4. The temperature Dantrolene may be stored at.
- 6. Heart rate does this during an MH crisis.
- 8. Once mixed, the contents of the dantrolene sodium vial should be administered within _____ hours.
- 10. The brand name for dantrolene sodium that turns orange upon mixing.
- 11. According to MHAUS, there are how many states that exhibit a high incidence of MH within the U.S.
- 14. Dantrolene must be available within _____ minutes of deciding to treat MH.

* Content reviewed by the CCI Nursing Education Department for alignment with clinical practice standards. CCI does not require, recommend, or endorse specific training programs in specialized practice areas for any of its exams. This is an example of a future points activity for recertification in collaboration with CCI.