


SURGIQUIZ



Breast Surgery CW138

To complete this activity, the participant will need to access and read Trotter, K.J., Neil, J.A. (2019). Breast Surgery. In J.C. Rothrock (Ed.). *Alexander's Care of the Patient in Surgery* (pp. 549 - 570). St. Louis, MO: Elsevier



Across

- 1. Also known as internal radiation.
- 4. Most common screening method for asymptomatic women.
- 5. More than 75% of lymph drainage drains into these lymph nodes.
- 6. The percentage of males affected with breast cancer.
- 8. The pigmented circular tissue surrounding the nipple.
- Class of genes known as tumor suppressors and when mutated are linked to hereditary breast cancers.
- 13. The fascia that breast tissue lies on.(2 words)

Down

- An alternative to needle localization for lesion identification that may be performed up to five days before surgery. (3 words)
- A breast conserving surgery that is also known as a partial mastectomy.
- 7. Most common benign breast lesion.
- This cancer in situ begins in the milk glands of the breast.
- 10. Sentinel lymph nodes are sent to pathology as a _____ specimen.
- 12. Vigorous scrubbing of the surgical site can result in possible _____ of cancer cells from the main mass.

^{*} Content reviewed by the CCI Nursing Education Department for alignment with clinical practice standards. CCI does not require, recommend, or endorse specific training programs in specialized practice areas for any of its exams. This is an example of a future points activity for recertification in collaboration with CCI.